

ON THE NUMBER OF SOLUTIONS TO THE ASYMPTOTIC PLATEAU PROBLEM

BARIS COSKUNUZER

ABSTRACT. We give a simple topological argument to show that the number of solutions to the asymptotic Plateau problem in hyperbolic space is generically unique. In particular, we show that the space of closed, codimension-1 submanifolds of $S_{\infty}^{n-1}(\mathbf{H}^n)$ which bound a unique absolutely area minimizing hypersurface in $\mathbf{H}^n$ is dense in the space of codimension-1 closed submanifolds of $S_{\infty}^{n-1}(\mathbf{H}^n)$. In dimension 3, we also prove that the set of uniqueness curves in $S_{\infty}^2(\mathbf{H}^3)$ for least area planes is generic in the set of Jordan curves in $S_{\infty}^2(\mathbf{H}^3)$. We also give some nonuniqueness results for dimension 3, too.

1. INTRODUCTION

The asymptotic Plateau problem in hyperbolic space asks the existence of an absolutely area minimizing hypersurface $\Sigma \subset \mathbf{H}^n$ asymptotic to given closed codimension-1 submanifold Γ in $S_{\infty}^{n-1}(\mathbf{H}^n)$. This problem is solved by Michael Anderson in his seminal paper [A1]. He proved the existence of a solution for any given closed submanifold in the sphere at infinity. Then, Hardt and Lin studied the asymptotic regularity of these solutions in [HL], [Li]. Lang generalized Anderson's methods to solve this problem in Gromov-Hadamard spaces in [L].

On the other hand, on the number of the absolutely area minimizing hypersurfaces for a given asymptotic boundary, there are a few results so far. In [A1], Anderson showed that if the given asymptotic boundary Γ bounds a convex domain in $S_{\infty}^{n-1}(\mathbf{H}^n)$, then there exists a unique absolutely area minimizing hypersurface in $\mathbf{H}^n$. Then, Hardt and Lin generalized this result to the closed codimension-1 submanifolds bounding star shaped domains in $S_{\infty}^{n-1}(\mathbf{H}^n)$ in [HL]. Recently, the author showed a generic uniqueness result in dimension 3 for least area planes in [Co1], [Co2]. For other results on asymptotic Plateau problem, see the survey article [Co5].

The author is supported by NSF Grant DMS-0603532, EU-FP7 Grant IRG-226062 and TUBITAK Grant 107T642.

In this paper, we will show that the space of closed codimension-1 submanifolds in $S_{\infty}^{n-1}(\mathbf{H}^n)$ bounding a unique absolutely area minimizing hypersurface in $\mathbf{H}^n$ is dense in the space of all closed codimension-1 submanifolds in $S_{\infty}^{n-1}(\mathbf{H}^n)$ by using a simple topological argument. On the other hand, in dimension 3, we show that the space of Jordan curves in $S_{\infty}^2(\mathbf{H}^3)$ bounding a unique area minimizing surface in $\mathbf{H}^3$ is generic in the space of Jordan curves in $S_{\infty}^2(\mathbf{H}^3)$. Similarly, we show that the same is true for the least area planes in $\mathbf{H}^3$, too.

The main results of the paper are the following. The first one is about least area planes in $\mathbf{H}^3$.

Theorem 3.3. Let A be the space of simple closed curves in $S_{\infty}^2(\mathbf{H}^3)$ and let $A' \subset A$ be the subspace containing the simple closed curves in $S_{\infty}^2(\mathbf{H}^3)$ bounding a unique least area plane in $\mathbf{H}^3$. Then, A' is generic in A , i.e. $A - A'$ is a set of first category.

The second result is about absolutely area minimizing hypersurfaces.

Theorem 4.4. Let B be the space of connected closed codimension-1 submanifolds of $S_{\infty}^{n-1}(\mathbf{H}^n)$, and let $B' \subset B$ be the subspace containing the closed submanifolds of $S_{\infty}^{n-1}(\mathbf{H}^n)$ bounding a unique absolutely area minimizing hypersurface in $\mathbf{H}^n$. Then B' is dense in B .

Indeed, this subspace of closed submanifolds in $S_{\infty}^{n-1}(\mathbf{H}^n)$ bounding a unique absolutely area minimizing hypersurface is not only dense, but also generic in some sense. See Remark 4.2. On the other hand, in dimension 3, by combination of these two theorems, we get the following corollary.

Corollary 4.5. Let A be the space of simple closed curves in $S_{\infty}^2(\mathbf{H}^3)$ and let $A' \subset A$ be the subspace containing the simple closed curves in $S_{\infty}^2(\mathbf{H}^3)$ bounding a unique absolutely area minimizing surface in $\mathbf{H}^3$. Then, A' is generic in A , i.e. $A - A'$ is a set of first category.

The short outline of the technique for generic uniqueness is the following: For simplicity, we will focus on the case of the least area planes in $\mathbf{H}^3$. Let Γ_0 be a simple closed curve in $S_{\infty}^2(\mathbf{H}^3)$. First, we will show that either there exists a unique least area plane Σ_0 in $\mathbf{H}^3$ with $\partial_{\infty}\Sigma_0 = \Gamma_0$, or there exist two *disjoint* least area planes Σ_0^+, Σ_0^- in $\mathbf{H}^3$ with $\partial_{\infty}\Sigma_0^{\pm} = \Gamma_0$. Now, take a small neighborhood $N(\Gamma_0) \subset S_{\infty}^2(\mathbf{H}^3)$ which is an annulus. Then foliate $N(\Gamma_0)$ by simple closed curves $\{\Gamma_t\}$ where $t \in (-\epsilon, \epsilon)$, i.e. $N(\Gamma_0) \simeq \Gamma \times (-\epsilon, \epsilon)$. By the above fact, for any Γ_t either there exists a unique least area plane Σ_t , or there are two least area planes $\Sigma_t^{\pm}$ disjoint from each other. Also, since these are least area planes, if they have disjoint asymptotic boundary, then they are disjoint by Meeks-Yau exchange roundoff trick. This means,

if $t_1 < t_2$, then Σ_{t_1} is disjoint and *below* from Σ_{t_2} in $\mathbf{H}^3$. Consider this collection of least area planes. Note that for curves Γ_t bounding more than one least area plane, we have a canonical region N_t in $\mathbf{H}^3$ between the disjoint least area planes $\Sigma_t^\pm$, see Figure 1.

Now, $N(\Gamma)$ separates $S_\infty^2(\mathbf{H}^3)$ into two parts, and take a geodesic $\beta \subset \mathbf{H}^3$ which is asymptotic to two points belongs to these two different parts. This geodesic is transverse to the collection of these least area planes asymptotic to the curves in $\{\Gamma_t\}$. Also, a finite segment of this geodesic intersects the entire collection. Let the length of this finite segment be C . Now, the idea is to consider the *thickness* of the neighborhoods N_t assigned to the asymptotic curves $\{\Gamma_t\}$. Let s_t be the length of the segment I_t of β between Σ_t^+ and Σ_t^- , which is the *width* of N_t assigned to Γ_t . Then, the curves Γ_t bounding more than one least area plane have positive width, and contributes to total thickness of the collection, and the curves bounding unique least area plane has 0 width and do not contribute to the total thickness. Since $\sum_{t \in (-\epsilon, \epsilon)} s_t < C$, the total thickness is finite. This implies for only countably many $t \in (-\epsilon, \epsilon)$, $s_t > 0$, i.e. Γ_t bounds more than one least area plane. For the remaining uncountably many $t \in (-\epsilon, \epsilon)$, $s_t = 0$, and there exists a unique least area plane for those t . This proves the space of Jordan curves of uniqueness is dense in the space of Jordan curves in $S_\infty^2(\mathbf{H}^3)$. Then, we will show this space is not only dense, but also generic.

We should note that this technique is quite general and it can be applied to many different settings of the Plateau problem (See Concluding Remarks).

On the other hand, after the above generic uniqueness results, it is a reasonable question whether all simple closed curves in $S_\infty^2(\mathbf{H}^3)$ bound a unique absolutely area minimizing surface or a unique least area plane. It is still not known whether all simple closed curves in $S_\infty^2(\mathbf{H}^3)$ have a unique solution to the asymptotic Plateau problem or not. The only known results about nonuniqueness also come from Anderson in [A2]. He constructs examples of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ bounding more than one complete *minimal surface* in $\mathbf{H}^3$. These examples are also area minimizing in their topological class. However, none of them are absolutely area minimizing, i.e. a solution to the asymptotic Plateau problem.

In this paper, we show the existence of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ with nonunique solution to the asymptotic Plateau problem. In other words, we show that there are examples of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ which are the asymptotic boundaries of more than one absolutely area minimizing surface.

Theorem 5.2. There exists a simple closed curve Γ in $S_\infty^2(\mathbf{H}^3)$ such that Γ bounds more than one absolutely area minimizing surface $\{\Sigma_i\}$ in $\mathbf{H}^3$, i.e. $\partial_\infty \Sigma_i = \Gamma$.

Also, by using similar ideas, we show nonuniqueness for least area planes case in dimension 3.

Theorem 5.3. There exists a simple closed curve Γ in $S_\infty^2(\mathbf{H}^3)$ such that Γ bounds more than one least area plane $\{P_i\}$ in $\mathbf{H}^3$, i.e. $\partial_\infty P_i = \Gamma$.

The organization of the paper is as follows: In the next section we will cover some basic results which will be used in the following sections. In section 3, we will show the genericity result for least area planes in $\mathbf{H}^3$. Then in section 4, we will show the results on absolutely area minimizing hypersurfaces in $\mathbf{H}^n$. In Section 5, we will prove the nonuniqueness results. Finally in section 6, we will have some concluding remarks.

1.1. Acknowledgements: I would like to thank Yair Minsky and Michael Anderson for very useful conversations.

2. PRELIMINARIES

In this section, we will overview the basic results which we will use in the following sections. For details on the notions and results in this section, see the survey article [Co5].

First, we will give the definitions of area minimizing surfaces. First set of the definitions are about compact surfaces and hypersurfaces. The second set of the definitions are their generalizations to the noncompact surfaces and hypersurfaces.

Definition 2.1. (Compact Case) A *least area disk* (area minimizing disk) is a disk which has the smallest area among the disks with the same boundary. An *absolutely area minimizing surface* is a surface which has the smallest area among all the surfaces (with no topological restriction) with the same boundary. An *absolutely area minimizing hypersurface* is a hypersurface which has the smallest volume among all hypersurfaces with the same boundary.

Definition 2.2. (Noncompact Case) A *least area plane* is a plane such that any compact subdisk in the plane is a least area disk. We will also call a complete noncompact surface as *absolutely area minimizing surface* if any compact subsurface is an absolutely area minimizing surface. Similarly, we will call a complete noncompact hypersurface as *absolutely area minimizing hypersurface*, if any compact part (codimension-0 submanifold with boundary) of the hypersurface is an absolutely area minimizing hypersurface.

Now, we will quote the basic results on asymptotic Plateau problem.

Lemma 2.1. [A1] *Let Γ be a codimension-1 closed submanifold of $S_\infty^{n-1}(\mathbf{H}^n)$. Then there exists a complete, absolutely area minimizing $n - 1$ -rectifiable current Σ in $\mathbf{H}^n$ with $\partial_\infty \Sigma = \Gamma$.*

Note that the rectifiable current here is indeed a smooth hypersurface of $\mathbf{H}^n$ except for a singular set of Hausdorff dimension at most $n - 8$ by the regularity result stated below. For convenience, we will call area minimizing codimension-1 rectifiable currents as area minimizing hypersurfaces throughout the paper.

Lemma 2.2. [A2] *Let Γ be a Jordan curve in $S_\infty^2(\mathbf{H}^3)$. Then, there exists a complete, least area plane Σ in $\mathbf{H}^3$ asymptotic to Γ .*

The following fact about interior regularity theory of geometric measure theory is well-known.

Lemma 2.3. [Fe] *Let Σ be a $(n - 1)$ -dimensional area minimizing rectifiable current. Then Σ is a smooth, embedded manifold in the interior except for a singular set of Hausdorff dimension at most $n - 8$.*

Finally, we will state a theorem about limits of sequences of least area planes. Here, the limit is the pointwise limit of the planes, and for each limit point, there is a disk containing the point in the limit set such that it is the limit of a sequence of subdisks in the planes [Ga, Lemma 3.3].

Lemma 2.4. [Ga] *Let $\{\Sigma_i\}$ be a sequence of least area planes in $\mathbf{H}^3$ with $\partial_\infty \Sigma_i = \Gamma_i \subset S_\infty^2(\mathbf{H}^3)$ simple closed curve for any i . If $\Gamma_i \rightarrow \Gamma$, then there exists a subsequence $\{\Sigma_{i_j}\}$ of $\{\Sigma_i\}$ such that $\Sigma_{i_j} \rightarrow \widehat{\Sigma}$ a collection of least area planes whose asymptotic boundaries are Γ .*

The next theorem is a similar limit theorem about absolutely area minimizing hypersurfaces in $\mathbf{H}^n$.

Lemma 2.5. *Let $\{\Gamma_i\}$ be a sequence of connected closed codimension-1 submanifolds in $S_\infty^{n-1}(\mathbf{H}^n)$ which are pairwise disjoint. Let $\{\Sigma_i\}$ be a sequence of complete absolutely area minimizing hypersurfaces in $\mathbf{H}^n$ with $\partial_\infty(\Sigma_i) = \Gamma_i$. If Γ_i converges to a closed codimension-1 submanifold Γ in $S_\infty^{n-1}(\mathbf{H}^n)$, then there exists a subsequence of $\{\Sigma_i\}$ which converges to a complete absolutely area minimizing hypersurfaces Σ in $\mathbf{H}^n$ with $\partial_\infty \Sigma = \Gamma$.*

Proof: By the proof of the Lemma 2.1, all we need to show that we can induce a suitable sequence of compact absolutely area minimizing hypersurfaces $\{S_i\}$ from $\{\Sigma_i\}$ where $\gamma_i = \partial S_i$ converges to Γ in $S_\infty^2(\mathbf{H}^3)$.

Let $K > 0$ be sufficiently large so that $\Sigma_i \cap N_K(CH(\Gamma)) \neq \emptyset$ where $N_K(CH(\Gamma))$ is the K neighborhood of the convex hull of Γ . Then, let $S_i = N_K(CH(\Gamma)) \cap \Sigma_i$.

Since $\Sigma_i \subset CH(\Gamma_i)$ and $\Gamma_i \rightarrow \Gamma$, then ∂S_i must converge to Γ asymptotically. Then, by the proof of Lemma 2.1 with slight modification for the uniform mass bounds coming from $N_K(CH(\Gamma))$, there exists a subsequence $\{S_{i_j}\}$ which converges to a complete absolutely area minimizing hypersurface Σ in $\mathbf{H}^n$ with $\partial_\infty \Sigma = \Gamma$. $\square$

3. LEAST AREA PLANES IN $\mathbf{H}^3$

In this section, we will prove that the space of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ bounding a unique least area plane in $\mathbf{H}^3$ is generic in the space of simple closed curves in $S_\infty^2(\mathbf{H}^3)$.

First, we will show that if two least area planes have disjoint asymptotic boundaries, then they are disjoint.

Lemma 3.1. *Let Γ_1 and Γ_2 be two disjoint simple closed curves in $S_\infty^2(\mathbf{H}^3)$. If Σ_1 and Σ_2 are least area planes in $\mathbf{H}^3$ with $\partial_\infty \Sigma_i = \Gamma_i$, then Σ_1 and Σ_2 are disjoint, too.*

Proof: Assume that $\Sigma_1 \cap \Sigma_2 \neq \emptyset$. Since asymptotic boundaries Γ_1 and Γ_2 are disjoint, the intersection cannot contain an infinite line. So, the intersection between Σ_1 and Σ_2 must contain a simple closed curve γ . Since Σ_1 and Σ_2 are also minimal, the intersection must be transverse on a subarc of γ by maximum principle.

Now, γ bounds two area minimizing disks D_1 and D_2 in $\mathbf{H}^3$, with $D_i \subset \Sigma_i$. Now, take a larger subdisk E_1 of Σ_1 containing D_1 , i.e. $D_1 \subset E_1 \subset \Sigma_1$. By definition, E_1 is also an area minimizing disk. Now, modify E_1 by swapping the disks D_1 and D_2 . Then, we get a new disk $E'_1 = \{E_1 - D_1\} \cup D_2$. Now, E_1 and E'_1 have same area, but E'_1 have a folding curve along γ . By smoothing out this curve as in [MY], we get a disk with smaller area, which contradicts to E_1 being area minimizing. Note that this technique is known as Meeks-Yau exchange roundoff trick. $\square$

The following lemma is very essential for our technique. Mainly, the lemma says that for any given simple closed curve Γ in $S_\infty^2(\mathbf{H}^3)$, either there exists a unique least area plane Σ in $\mathbf{H}^3$ asymptotic to Γ , or there exist two least area planes $\Sigma^\pm$ in $\mathbf{H}^3$ which are asymptotic to Γ and disjoint from each other. Even though this lemma is also proven in [Co4], because of its importance for the technique, and to set the notation for the main result, we give a proof here. Note that Brian White proved a similar version of this lemma by using geometric measure theory methods in [Wh1].

Lemma 3.2. *Let Γ be a simple closed curve in $S_\infty^2(\mathbf{H}^3)$. Then either there exists a unique least area plane Σ in $\mathbf{H}^3$ with $\partial_\infty \Sigma = \Gamma$, or there are two canonical disjoint extremal least area planes Σ^+ and Σ^- in $\mathbf{H}^3$ with*

$\partial_\infty \Sigma^\pm = \Gamma$. Moreover, any least area plane Σ' with $\partial_\infty \Sigma' = \Gamma$ is disjoint from $\Sigma^\pm$, and it is captured in the region bounded by Σ^+ and Σ^- in $\mathbf{H}^3$.

Proof: Let Γ be a simple closed curve in $S_\infty^2(\mathbf{H}^3)$. Γ separates $S_\infty^2(\mathbf{H}^3)$ into two parts, say D^+ and D^- . Define sequences of pairwise disjoint simple closed curves $\{\Gamma_i^+\}$ and $\{\Gamma_i^-\}$ such that $\Gamma_i^+ \subset D^+$, and $\Gamma_i^- \subset D^-$ for any i , and $\Gamma_i^+ \rightarrow \Gamma$, and $\Gamma_i^- \rightarrow \Gamma$.

By Lemma 2.2, for any $\Gamma_i^+ \subset S_\infty^2(\mathbf{H}^3)$, there exists a least area plane Σ_i^+ in $\mathbf{H}^3$ asymptotic to Γ_i^+ . This defines a sequence of least area planes $\{\Sigma_i^+\}$. Now, by using Lemma 2.4, we take the limit of a convergent subsequence. In the limit we get a collection of least area planes $\widehat{\Sigma}^+$ with $\partial_\infty \widehat{\Sigma}^+ = \Gamma$, as $\partial_\infty \Sigma_i^+ = \Gamma_i^+ \rightarrow \Gamma$.

Now, we claim that the collection $\widehat{\Sigma}^+$ consists of only one least area plane. Assume that there are two least area planes Σ_a^+ and Σ_b^+ in the collection $\widehat{\Sigma}^+$. Since $\partial_\infty \Sigma_a^+ = \partial_\infty \Sigma_b^+ = \Gamma$, Σ_a^+ and Σ_b^+ might not be disjoint, but they are disjoint from least area planes in the sequence, i.e. $\Sigma_i^+ \cap \Sigma_{a,b}^+ = \emptyset$ for any i , by Lemma 3.1.

If Σ_a^+ and Σ_b^+ are disjoint, say Σ_a^+ is *above* Σ_b^+ . By Lemma 3.1, we know that for any i , Σ_i^+ is *above* both Σ_a^+ and Σ_b^+ . However this means that Σ_a^+ is a barrier between the sequence $\{\Sigma_i^+\}$ and Σ_b^+ , and so, Σ_b^+ cannot be limit of this sequence, which is a contradiction.

If Σ_a^+ and Σ_b^+ are not disjoint, then they intersect each other, and in some region, Σ_b^+ is *above* Σ_a^+ . However since Σ_a^+ is the limit of the sequence $\{\Sigma_i^+\}$, this would imply Σ_b^+ must intersect planes Σ_i^+ for sufficiently large i . However, this contradicts the fact that Σ_b^+ is disjoint from Σ_i^+ for any i , as they have disjoint asymptotic boundary. So, there exists a unique least area plane Σ^+ in the collection $\widehat{\Sigma}^+$. Similarly, $\widehat{\Sigma}^- = \Sigma^-$. By using similar arguments, one can conclude that these least area planes Σ^+ , and Σ^- are canonical, i.e. independent of the choice of the sequence $\{\Gamma_i^+\}$ and $\{\Gamma_i^-\}$.

Now, let Σ' be any least area plane with $\partial_\infty \Sigma' = \Gamma$. If $\Sigma' \cap \Sigma^+ \neq \emptyset$, then some part of Σ' must be *above* Σ^+ . Since $\Sigma^+ = \lim \Sigma_i^+$, for sufficiently large i , $\Sigma' \cap \Sigma_i^+ \neq \emptyset$. However, $\partial_\infty \Sigma_i^+ = \Gamma_i^+$ is disjoint from $\Gamma = \partial_\infty \Sigma'$. Then, by Lemma 3.1, Σ' must be disjoint from Σ_i^+ . This is a contradiction.

Similarly, this is true for Σ^- , too. Moreover, let $N \subset \mathbf{H}^3$ be the region between Σ^+ and Σ^- , i.e. $\partial N = \Sigma^+ \cup \Sigma^-$. Then by construction, N is also a canonical region for Γ , and for any least area plane Σ' with $\partial_\infty \Sigma' = \Gamma$, Σ' is contained in the region N , i.e. $\Sigma' \subset N$. This shows that if $\Sigma^+ = \Sigma^-$, there exists a unique least area plane asymptotic to Γ . If $\Sigma^+ \neq \Sigma^-$, then they must be disjoint. $\square$

Now, we are going to prove the main theorem of the section. This theorem says that for a generic simple closed curve in $S_\infty^2(\mathbf{H}^3)$, there exists a unique least area plane.

Theorem 3.3. *Let A be the space of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ and let $A' \subset A$ be the subspace containing the simple closed curves in $S_\infty^2(\mathbf{H}^3)$ bounding a unique least area plane in $\mathbf{H}^3$. Then, A' is generic in A , i.e. $A - A'$ is a set of first category.*

Proof: We will prove this theorem in 2 steps.

Claim 1: A' is dense in A as a subspace of $C^0(S^1, S_\infty^2(\mathbf{H}^3))$ with the supremum metric.

Proof: A is the space of Jordan curves in $S_\infty^2(\mathbf{H}^3)$. Then, $A = \{\alpha \in C^0(S^1, S^2) \mid \alpha(S^1) \text{ is an embedding}\}$.

Now, let $\Gamma_0 \in A$ be a simple closed curve in $S_\infty^2(\mathbf{H}^3)$. Since Γ_0 is simple, there exists a small neighborhood $N(\Gamma_0)$ of Γ_0 which is an annulus in $S_\infty^2(\mathbf{H}^3)$. Let $\Gamma : (-\epsilon, \epsilon) \rightarrow A$ be a small path in A through Γ_0 such that $\Gamma(t) = \Gamma_t$ and $\{\Gamma_t\}$ foliates $N(\Gamma)$ with simple closed curves Γ_t . In other words, $\{\Gamma_t\}$ are pairwise disjoint simple closed curves, and $N(\Gamma_0) = \bigcup_{t \in (-\epsilon, \epsilon)} \Gamma_t$.

Now, $N(\Gamma_0)$ separates $S_\infty^2(\mathbf{H}^3)$ into two parts, say D^+ and D^- , i.e. $S_\infty^2(\mathbf{H}^3) = N(\Gamma_0) \cup D^+ \cup D^-$. Let p^+ be a point in D^+ and let p^- be a point in D^- such that for a small δ , $B_\delta(p^\pm)$ are in the interior of $D^\pm$. Let β be the geodesic in $\mathbf{H}^3$ asymptotic to p^+ and p^- .

By Lemma 3.2, for any Γ_t either there exists a unique least area plane Σ_t in $\mathbf{H}^3$, or there is a canonical region N_t in $\mathbf{H}^3$ between the canonical least area planes Σ_t^+ and Σ_t^- (In Figure 1, Γ_t and Γ_s bound more than one least area plane in $\mathbf{H}^3$, whereas Γ_0 bounds a unique least area plane Σ_0 in $\mathbf{H}^3$). With abuse of notation, if Γ_t bounds a unique least area plane Σ_t in $\mathbf{H}^3$, define $N_t = \Sigma_t$ as a degenerate canonical neighborhood for Γ_t (In Figure 1, N_t and N_s represent nondegenerate canonical neighborhoods, and $N_0 = \Sigma_0$ represents degenerate canonical neighborhood.). Then, let $\hat{N} = \{N_t\}$ be the collection of these degenerate and nondegenerate canonical neighborhoods for $t \in (-\epsilon, \epsilon)$. Clearly, degenerate neighborhood N_t means Γ_t bounds unique least area plane, and nondegenerate neighborhood N_s means that Γ_s bounds more than one least area plane. Note that by Lemma 3.1, all canonical neighborhoods in the collection are pairwise disjoint. On the other hand, by construction the geodesic β intersects all the canonical neighborhoods in the collection $\hat{N}$.

We claim that the part of β which intersects $\hat{N}$ is a finite line segment. Let P^+ be the geodesic plane asymptotic to round circle $\partial B_\delta(p^+)$ in D^+ .




FIGURE 1. A finite segment of geodesic γ intersects the collection of least area planes Σ_t in $\mathbf{H}^n$ asymptotic to Γ_t in $S_{\infty}^{n-1}(\mathbf{H}^n)$.

Similarly, define P^- . By Lemma 3.1, $P^\pm$ are disjoint from the collection of canonical regions $\widehat{N}$. Let $\beta \cap P^\pm = \{q^\pm\}$. Then the part of β which intersects $\widehat{N}$ is the line segment $l \subset \beta$ with endpoints q^+ and q^- . Let C be the length of this line segment l .

Now, for each $t \in (-\epsilon, \epsilon)$, we will assign a real number $s_t \geq 0$. If there exists a unique least area plane Σ_t in $\mathbf{H}^3$ for Γ_t (N_t is degenerate), then let s_t be 0. If not, let $I_t = \beta \cap N_t$, and s_t be the length of I_t . Clearly if Γ_t bounds more than one least area plane (N_t is nondegenerate), then $s_t > 0$. Also, it is clear that for any t , $I_t \subset l$ and $I_t \cap I_s = \emptyset$ for any $t \neq s$. Then, $\sum_{t \in (-\epsilon, \epsilon)} s_t < C$ where C is the length of l . This means for only countably many $t \in (-\epsilon, \epsilon)$, $s_t > 0$. So, there are only countably many nondegenerate N_t for $t \in (-\epsilon, \epsilon)$. Hence, for all other t , N_t is degenerate. This means there exist uncountably many $t \in (-\epsilon, \epsilon)$, where Γ_t bounds a unique least area plane. Since Γ_0 is arbitrary, this proves A' is dense in A . $\square$

Claim 2: A' is generic in A , i.e. $A - A'$ is a set of first category.

Proof: We will prove that A' is countable intersection of open dense subsets of a complete metric space. Then the result will follow by Baire category theorem.

Since the space of continuous maps from circle to sphere $C^0(S^1, S^2)$ is complete with supremum metric, then the closure of A in $C^0(S^1, S^2)$, $\bar{A} \subset C^0(S^1, S^2)$, is also complete.

Now, we will define a sequence of open dense subsets $U^i \subset A$ such that their intersection will give us A' . Let $\Gamma \in A$ be a simple closed curve in $S_\infty^2(\mathbf{H}^3)$, as in the Claim 1. Let $N(\Gamma) \subset S_\infty^2(\mathbf{H}^3)$ be a neighborhood of Γ in $S_\infty^2(\mathbf{H}^3)$, which is an open annulus. Then, define an open neighborhood U_Γ of Γ in A , such that $U_\Gamma = \{\alpha \in A \mid \alpha(S^1) \subset N(\Gamma), \alpha \text{ is homotopic to } \Gamma\}$. Clearly, $A = \bigcup_{\Gamma \in A} U_\Gamma$. Now, define a geodesic β_Γ as in Claim 1, which intersects all the least area planes asymptotic to curves in U_Γ .

Now, for any $\alpha \in U_\Gamma$, by Lemma 3.2, there exists a canonical region N_α in $\mathbf{H}^3$ (which can be degenerate if α bounds a unique least area plane). Let $I_{\alpha, \Gamma} = N_\alpha \cap \beta_\Gamma$. Then let $s_{\alpha, \Gamma}$ be the length of $I_{\alpha, \Gamma}$ ($s_{\alpha, \Gamma}$ is 0 if N_α degenerate). Hence, for every element α in U_Γ , we assign a real number $s_{\alpha, \Gamma} \geq 0$.

Now, we define the sequence of open dense subsets in U_Γ . Let $U_\Gamma^i = \{\alpha \in U_\Gamma \mid s_{\alpha, \Gamma} < 1/i\}$. We claim that U_Γ^i is an open subset of U_Γ and A . Let $\alpha \in U_\Gamma^i$, and let $s_{\alpha, \Gamma} = \lambda < 1/i$. So, the interval $I_{\alpha, \Gamma} \subset \beta_\Gamma$ has length λ . let $I' \subset \beta_\Gamma$ be an interval containing $I_{\alpha, \Gamma}$ in its interior, and has length less than $1/i$. By the proof of Claim 1, we can find two simple closed curves $\alpha^+, \alpha^- \in U_\Gamma$ with the following properties.

- $\alpha^\pm$ are disjoint from α ,
- $\alpha^\pm$ are lying in opposite sides of α in $S_\infty^2(\mathbf{H}^3)$,
- $\alpha^\pm$ bounds unique least area planes $\Sigma_{\alpha^\pm}$,
- $\Sigma_{\alpha^\pm} \cap \beta_\Gamma \subset I'$.

The existence of such curves is clear from the proof Claim 1, as if one takes any foliation $\{\alpha_t\}$ of a small neighborhood of α in $S_\infty^2(\mathbf{H}^3)$, there are uncountably many curves in the family bounding a unique least area plane, and one can choose sufficiently close pair of curves to α , to ensure the conditions above.

After finding $\alpha^\pm$, consider the open annulus F_α in $S_\infty^2(\mathbf{H}^3)$ bounded by α^+ and α^- . Let $V_\alpha = \{\gamma \in U_\Gamma \mid \gamma(S^1) \subset F_\alpha, \gamma \text{ is homotopic to } \alpha\}$. Clearly, V_α is an open subset of U_Γ . If we can show $V_\alpha \subset U_\Gamma^i$, then this proves U_Γ^i is open for any i and any $\Gamma \in A$.

Let $\gamma \in V_\alpha$ be any curve, and N_γ be its canonical neighborhood given by Lemma 3.2. Since $\gamma(S^1) \subset F_\alpha$, α^+ and α^- lie in opposite sides of γ in $S_\infty^2(\mathbf{H}^3)$. This means Σ_{α^+} and Σ_{α^-} lie in opposite sides of N_γ . By choice

of $\alpha^\pm$, this implies $N_\gamma \cap \beta_\Gamma = I_{\gamma,\Gamma} \subset I'$. So, the length $s_{\gamma,\Gamma}$ is less than $1/i$. This implies $\gamma \in U_\Gamma^i$, and so $V_\alpha \subset U_\Gamma^i$. Hence, U_Γ^i is open in U_Γ and A .

Now, we can define the sequence of open dense subsets. let $U^i = \bigcup_{\Gamma \in A} U_\Gamma^i$ be an open subset of A . Since, the elements in A' represent the curves bounding a unique least area plane, for any $\alpha \in A'$, and for any $\Gamma \in A$, $s_{\alpha,\Gamma} = 0$. This means $A' \subset U^i$ for any i . By Claim 1, U^i is open dense in A for any $i > 0$.

As we mention at the beginning of the proof, since the space of continuous maps from circle to sphere $C^0(S^1, S^2)$ is complete with supremum metric, then the closure $\bar{A}$ of A in $C^0(S^1, S^2)$ is also complete metric space. Since A' is dense in A , it is also dense in $\bar{A}$. As A is open in $C^0(S^1, S^2)$, this implies U^i is a sequence of open dense subsets of $\bar{A}$. On the other hand, since $s_{\alpha,\Gamma} = 0$ for any $\alpha \in A'$, and for any $\Gamma \in A$, $A' = \bigcap_{i>0} U^i$. Then, $A - A'$ is a set of first category, by Baire Category Theorem. Hence, A' is generic in A . $\square$

$\square$

Remark 3.1. This result is similar to the generic uniqueness result in [Co2]. In [Co2], we used a heavy machinery of analysis to prove that there exists an open dense subset in the space of $C^{3,\mu}$ -smooth embeddings of circle into sphere, where any simple closed curve in this space bounds a unique least area plane in $\mathbf{H}^3$. In the above result, the argument is fairly simple, and does not use the analytical machinery.

4. AREA MINIMIZING HYPERSURFACES IN $\mathbf{H}^n$

In this section, we will show that the space of codimension-1 closed submanifolds of $S_\infty^{n-1}(\mathbf{H}^n)$ bounding a unique absolutely area minimizing hypersurface in $\mathbf{H}^n$ is dense in the space of all codimension-1 closed submanifolds of $S_\infty^{n-1}(\mathbf{H}^n)$. Indeed, we will show that in some sense, a generic closed manifold Γ in $S_\infty^{n-1}(\mathbf{H}^n)$ bounds a unique absolutely area minimizing hypersurface Σ in $\mathbf{H}^n$. The idea is similar to the previous section.

First, we need to show a simple topological lemma.

Lemma 4.1. *Any codimension-1 closed submanifold Γ of S^{n-1} is separating. Moreover, if Σ is a hypersurface with boundary in the closed unit ball B^n , where the boundary $\partial\Sigma \subset \partial B^n$, then Σ is also separating, too.*

Proof: A non-separating codimension-1 closed submanifold in S^{n-1} gives a nontrivial homology in $n - 2$ level. However since $H_{n-2}(S^{n-1})$ is trivial, this is a contradiction.




FIGURE 2. S_1 is the part of Σ_1 lying below Σ_2 , and S_2 is the part of Σ_2 lying above Σ_1 . After swaping S_1 and S_2 , we get a new area minimizing hypersurface Σ'_1 with singularity along $\alpha = \Sigma_1 \cap \Sigma_2$.

Let Σ be as in the assumption. Take the double of B^n , then $B^n \sqcup \widehat{B}^n = S^n$, and $\Sigma \sqcup \widehat{\Sigma}$ is a codimension-1 closed submanifold of S^n . By above, $\Sigma \sqcup \widehat{\Sigma}$ is separating in S^n . Hence, Σ is separating in B^n . $\square$

Now, we will prove a disjointness lemma analogous to Lemma 3.1. This lemma roughly says that if asymptotic boundaries of two absolutely area minimizing hypersurfaces in $\mathbf{H}^n$ are disjoint in $S_\infty^{n-1}(\mathbf{H}^n)$, then they are disjoint in $\mathbf{H}^n$.

Lemma 4.2. *Let Γ_1 and Γ_2 be two disjoint connected closed codimension-1 submanifolds in $S_\infty^{n-1}(\mathbf{H}^n)$. If Σ_1 and Σ_2 are absolutely area minimizing hypersurfaces in $\mathbf{H}^n$ with $\partial_\infty \Sigma_i = \Gamma_i$, then Σ_1 and Σ_2 are disjoint, too.*

Proof: Assume that the absolutely area minimizing hypersurfaces are not disjoint, i.e. $\Sigma_1 \cap \Sigma_2 \neq \emptyset$. By Lemma 4.1, Σ_1 and Σ_2 separates $\mathbf{H}^n$ into two parts. So, say $\mathbf{H}^n - \Sigma_i = \Omega_i^+ \cup \Omega_i^-$.

Now, consider the intersection of hypersurfaces $\alpha = \Sigma_1 \cap \Sigma_2$. Since the asymptotic boundaries Γ_1 and Γ_2 are disjoint in $S_\infty^{n-1}(\mathbf{H}^n)$, by using the regularity results of Hardt and Lin in [HL], we can conclude that the intersection set α is in compact part of $\mathbf{H}^n$. Moreover, by maximum principle [Si], the intersection cannot have isolated tangential intersections.

Now, without loss of generality, we assume that Σ_1 is above Σ_2 (the non-compact part of Σ_1 lies in Ω_2^+). Now define the compact subhypersurfaces S_i in Σ_i as $S_1 = \Sigma_1 \cap \Omega_2^-$, and $S_2 = \Sigma_2 \cap \Omega_1^+$. In other words, S_1 is the part of Σ_1 lying below Σ_2 , and S_2 is the part of Σ_2 lying above Σ_1 . Then, $\partial S_1 = \partial S_2 = \alpha$. See Figure 2.

On the other hand, since Σ_1 and Σ_2 are absolutely area minimizing, then by definition, so are S_1 and S_2 , too. Then by swaping the surfaces, we can get new absolutely area minimizing hypersurfaces. In other words, let $\Sigma'_1 = \{\Sigma_1 - S_1\} \cup S_2$, and $\Sigma'_2 = \{\Sigma_2 - S_2\} \cup S_1$ are also absolutely area minimizing hypersurfaces. However, in this new hypersurfaces, we will have a singularity set along α , which contradicts to regularity theorem for absolutely area minimizing hypersurfaces, i.e. Lemma 2.3. $\square$

Lemma 4.3. *Let Γ be a connected closed codimension-1 submanifold of $S_{\infty}^{n-1}(\mathbf{H}^n)$. Then either there exists a unique absolutely area minimizing hypersurface Σ in $\mathbf{H}^n$ asymptotic to Γ , or there are two canonical disjoint extremal absolutely area minimizing hypersurfaces Σ^+ and Σ^- in $\mathbf{H}^n$ asymptotic to Γ .*

Proof: Let Γ be a connected closed codimension-1 submanifold of $S_{\infty}^{n-1}(\mathbf{H}^n)$. Then by Lemma 4.1, Γ separates $S_{\infty}^{n-1}(\mathbf{H}^n)$ into two parts, say Ω^+ and Ω^- . Define sequences of pairwise disjoint closed submanifolds of the same topological type $\{\Gamma_i^+\}$ and $\{\Gamma_i^-\}$ in $S_{\infty}^{n-1}(\mathbf{H}^n)$ such that $\Gamma_i^+ \subset \Omega^+$, and $\Gamma_i^- \subset \Omega^-$ for any i , and $\Gamma_i^+ \rightarrow \Gamma$, and $\Gamma_i^- \rightarrow \Gamma$ in Hausdorff metric. In other words, $\{\Gamma_i^+\}$ and $\{\Gamma_i^-\}$ converges to Γ from opposite sides.

By Lemma 2.1, for any $\Gamma_i^+ \subset S_{\infty}^2(\mathbf{H}^3)$, there exists an absolutely area minimizing hypersurface Σ_i^+ in $\mathbf{H}^n$. This defines a sequence of absolutely area minimizing hypersurfaces $\{\Sigma_i^+\}$. By Lemma 2.5, we get a convergent subsequence $\Sigma_{i_j}^+ \rightarrow \Sigma^+$. Hence, we get the absolutely area minimizing hypersurface Σ^+ in $\mathbf{H}^n$ asymptotic to Γ . Similarly, we get the absolutely area minimizing hypersurface Σ^- in $\mathbf{H}^n$ asymptotic to Γ . Similar arguments show that these absolutely area minimizing hypersurfaces $\Sigma^{\pm}$ are canonical by their construction, i.e. independent of the choice of the sequence $\{\Gamma_i^{\pm}\}$ and $\{\Sigma_i^{\pm}\}$.

Assume that $\Sigma^+ \neq \Sigma^-$, and they are not disjoint. Since these are absolutely area minimizing hypersurfaces, nontrivial intersection implies some part of Σ^- lies *above* Σ^+ . Since $\Sigma^+ = \lim \Sigma_{i_j}^+$, Σ^- must also intersect some $\Sigma_{i_j}^+$ for sufficiently large i_j . However by Lemma 4.2, $\Sigma_{i_j}^+$ is disjoint from Σ^- as $\partial_{\infty} \Sigma_{i_j}^+ = \Gamma_{i_j}^+$ is disjoint from $\partial_{\infty} \Sigma^- = \Gamma$. This is a contradiction. This shows Σ^+ and Σ^- are disjoint.

Similar arguments show that $\Sigma^{\pm}$ are disjoint from any absolutely area minimizing hypersurface Σ' asymptotic Γ . As the sequences of Σ_i^+ and Σ_i^- forms a barrier for other absolutely area minimizing hypersurfaces asymptotic to Γ , any such absolutely area minimizing hypersurface must lie in the region bounded by Σ^+ and Σ^- in $\mathbf{H}^n$. This shows that if $\Sigma^+ = \Sigma^-$, then there exists a unique absolutely area minimizing hypersurface asymptotic to Γ . $\square$

Remark 4.1. By above theorem and its proof, if Γ bounds more than one absolutely area minimizing hypersurface, then there exists a canonical region N_Γ in $\mathbf{H}^n$ asymptotic to Γ such that N_Γ is the region between the canonical absolutely area minimizing hypersurfaces Σ^+ and Σ^- . Moreover, by using similar ideas to the proof of Lemma 3.2, one can show that any absolutely area minimizing hypersurface in $\mathbf{H}^n$ asymptotic to Γ is in the region N_Γ .

Now, we can prove the main result of the paper.

Theorem 4.4. *Let B be the space of connected closed codimension-1 submanifolds of $S_\infty^{n-1}(\mathbf{H}^n)$, and let $B' \subset B$ be the subspace containing the closed submanifolds of $S_\infty^{n-1}(\mathbf{H}^n)$ bounding a unique absolutely area minimizing hypersurface in $\mathbf{H}^n$. Then B' is dense in B .*

Proof: Let B be the space of connected closed codimension-1 submanifolds of $S_\infty^{n-1}(\mathbf{H}^n)$ with Hausdorff metric. Let $\Gamma_0 \in B$ be a closed submanifold in $S_\infty^{n-1}(\mathbf{H}^n)$. Since Γ_0 is closed submanifold, there exists a small regular neighborhood $N(\Gamma_0)$ of Γ_0 in $S_\infty^{n-1}(\mathbf{H}^n)$, which is homeomorphic to $\Gamma_0 \times I$. Let $\Gamma : (-\epsilon, \epsilon) \rightarrow B$ be a small path in B through Γ_0 such that $\Gamma(t) = \Gamma_t$ and $\{\Gamma_t\}$ foliates $N(\Gamma_0)$ with closed submanifolds homeomorphic to Γ_0 . In other words, $\{\Gamma_t\}$ are pairwise disjoint closed submanifolds homeomorphic to Γ_0 , and $N(\Gamma_0) = \bigcup_{t \in (-\epsilon, \epsilon)} \Gamma_t$.

By Lemma 4.1, $N(\Gamma_0)$ separates $S_\infty^{n-1}(\mathbf{H}^n)$ into two parts, say Ω^+ and Ω^- , i.e. $S_\infty^{n-1}(\mathbf{H}^n) = N(\Gamma_0) \cup \Omega^+ \cup \Omega^-$. Let p^+ be a point in Ω^+ and let p^- be a point in Ω^- such that for a small δ , $B_\delta(p^\pm)$ are in the interior of $\Omega^\pm$. Let β be the geodesic in $\mathbf{H}^n$ asymptotic to p^+ and p^- .

By Lemma 4.3 and Remark 4.1, for any Γ_t either there exists a unique absolutely area minimizing hypersurface Σ_t in $\mathbf{H}^n$, or there is a canonical region N_t in $\mathbf{H}^n$ asymptotic to Γ_t , namely the region between the canonical absolutely area minimizing hypersurfaces Σ_t^+ and Σ_t^- . With abuse of notation, if Γ_t bounds a unique absolutely area minimizing hypersurface Σ_t in $\mathbf{H}^n$, define $N_t = \Sigma_t$ as a degenerate canonical neighborhood for Γ_t . Then, let $\hat{N} = \{N_t\}$ be the collection of these degenerate and nondegenerate canonical neighborhoods for $t \in (-\epsilon, \epsilon)$. Clearly, degenerate neighborhood N_t means Γ_t bounds a unique absolutely area minimizing hypersurface, and nondegenerate neighborhood N_s means that Γ_s bounds more than one absolutely area minimizing hypersurfaces. Note that by Lemma 4.2, all canonical neighborhoods in the collection are pairwise disjoint. On the other hand, by Lemma 4.1, the geodesic β intersects all the canonical neighborhoods in the collection $\hat{N}$.

We claim that the part of β which intersects $\hat{N}$ is a finite line segment. Let P^+ be the geodesic hyperplane asymptotic to round sphere $\partial B_\delta(p^+)$ in Ω^+ . Similarly, define P^- . By Lemma 4.2, $P^\pm$ are disjoint from the collection

of canonical regions $\widehat{N}$. Let $\beta \cap P^\pm = \{q^\pm\}$. Then the part of β which intersects $\widehat{N}$ is the line segment $l \subset \beta$ with endpoints q^+ and q^- . Let C be the length of this line segment l .

Now, for each $t \in (-\epsilon, \epsilon)$, we will assign a real number $s_t \geq 0$. If there exists a unique absolutely area minimizing hypersurface Σ_t for Γ_t (N_t is degenerate), then let s_t be 0. If not, let $I_t = \beta \cap N_t$, and s_t be the length of I_t . Clearly if Γ_t bounds more than one least area plane (N_t is nondegenerate), then $s_t > 0$. Also, it is clear that for any t , $I_t \subset l$ and $I_t \cap I_s = \emptyset$ for any $t \neq s$. Then, $\sum_{t \in (-\epsilon, \epsilon)} s_t < C$ where C is the length of l . This means for only countably many $t \in (-\epsilon, \epsilon)$, $s_t > 0$. So, there are only countably many nondegenerate N_t for $t \in (-\epsilon, \epsilon)$. Hence, for all other t , N_t is degenerate. This means there exist uncountably many $t \in (-\epsilon, \epsilon)$, where Γ_t bounds a unique absolutely area minimizing hypersurface. Since Γ_0 is arbitrary, this proves B' is dense in B . $\square$

Remark 4.2. This density result can be generalized to a genericity in some sense. Here, if one does not consider the whole space of closed codimension-1 submanifolds of $S_\infty^{n-1}(\mathbf{H}^n)$, but specify the topological type of the closed submanifolds, by using similar arguments to the proof of Claim 2 in Theorem 3.3, one can get a genericity result in that space. In other words, one can stratify the whole space of codimension-1 closed submanifolds by topological type, and get the genericity result in each strata.

Corollary 4.5. *Let A be the space of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ and let $A' \subset A$ be the subspace containing the simple closed curves in $S_\infty^2(\mathbf{H}^3)$ bounding a unique absolutely area minimizing surface in $\mathbf{H}^3$. Then, A' is generic in A , i.e. $A - A'$ is a set of first category.*

Proof: By Theorem 4.4, we know that A' is dense in A . Then by using the proof of Claim 2 in Theorem 3.3 in this setting, it is clear that A' is generic in A as a subspace of $C^0(S^1, S^2)$ with supremum metric. In other words, $A - A'$ is a set of first category. $\square$

Remark 4.3. This corollary shows that like in the case of least area planes (Theorem 3.3), a generic simple closed curve in $S_\infty^2(\mathbf{H}^3)$ also bounds a unique absolutely area minimizing surface.

5. NONUNIQUENESS RESULTS

In this section, we will show that there exists a simple closed curve in $S_\infty^2(\mathbf{H}^3)$ which is the asymptotic boundary of more than one absolutely area minimizing surface in $\mathbf{H}^3$. First, we need a lemma about the limits of absolutely area minimizing surfaces.

A short outline of the method is the following. We first construct a simple closed curve Γ_0 in $S_\infty^2(\mathbf{H}^3)$ which is the asymptotic boundary of more than one *minimal surface* in $\mathbf{H}^3$. Then, we foliate $S_\infty^2(\mathbf{H}^3)$ with simple closed curves $\{\Gamma_t\}$ where Γ_0 is a leaf in the foliation. We show that if each Γ_t bounds a unique absolutely area minimizing surface Σ_t in $\mathbf{H}^3$, then the family of surfaces $\{\Sigma_t\}$ must foliate the whole $\mathbf{H}^3$. However, since we chose Γ_0 to bound more than one minimal surface in $\mathbf{H}^3$, one of the surfaces must have a tangential intersection with one of the leaves in the foliation. This contradicts to the maximum principle for minimal surfaces.

Now, we quote a result on the existence of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ which are the asymptotic boundaries of more than one minimal surface in $\mathbf{H}^3$.

Lemma 5.1. [A2] *There is a set Δ of Jordan curves in $S_\infty^2(\mathbf{H}^3)$ such that for any $\Gamma \in \Delta$, there exist infinitely many complete, smoothly embedded minimal surfaces in $\mathbf{H}^3$ asymptotic to Γ .*

Remark 5.1. Alternatively, one can construct simple closed curves in $S_\infty^2(\mathbf{H}^3)$ bounding more than one minimal surface in $\mathbf{H}^3$ as follows. By using the technique in [Ha], one can construct a simple closed curve Γ in $S_\infty^2(\mathbf{H}^3)$ such that the absolutely area minimizing surface Σ asymptotic to Γ has positive genus (See Figure 3). Then, Σ separates $\mathbf{H}^3$ into two parts Ω^+ and Ω^- which are both mean convex domains. Then by using Meeks-Yau's results in [MY], one can get sequences of least area disks $\{D_i^\pm\}$ in $\Omega^\pm$ whose boundaries converges to Γ in $S_\infty^2(\mathbf{H}^3)$. By taking the limit, one can get two




FIGURE 3. Γ is a simple closed curve in $S_\infty^2(\mathbf{H}^3)$. γ_i^+ and γ_i^- are round circles in $S_\infty^2(\mathbf{H}^3)$ bounding the geodesic planes P_i^+ and P_i^- in $\mathbf{H}^3$.

least area planes to Γ . We used Σ as barrier to get distinct limit from the sequences $\{D_i^+\}$ and $\{D_i^-\}$. Note that the planes $M^\pm$ are least area just in $\Omega^\pm$. So, $M^\pm$ may not be least area in $\mathbf{H}^3$, but of course, they are still minimal planes.

Theorem 5.2. *There exists a simple closed curve Γ in $S_\infty^2(\mathbf{H}^3)$ such that Γ bounds more than one absolutely area minimizing surface $\{\Sigma_i\}$ in $\mathbf{H}^3$, i.e. $\partial_\infty \Sigma_i = \Gamma$.*

Proof: Assume that for any simple closed curve Γ in $S_\infty^2(\mathbf{H}^3)$, there exists a unique complete absolutely area minimizing surface Σ in $\mathbf{H}^3$ with $\partial_\infty \Sigma = \Gamma$. Let Γ_0 be a simple closed curve in $S_\infty^2(\mathbf{H}^3)$ such that M_1 and M_2 are two distinct minimal surfaces in $\mathbf{H}^3$ asymptotic to Γ as in Lemma 5.1.

Now, foliate $S_\infty^2(\mathbf{H}^3)$ by simple closed curves $\{\Gamma_t\}$ where Γ_0 is a leave in the foliation. Note that, there are only two singular leaves in the foliation which are points, and all other leaves are embedded simple closed curves in $S_\infty^2(\mathbf{H}^3)$. By assumption, for any Γ_t , there exists a unique absolutely area minimizing surface Σ_t in $\mathbf{H}^3$.

We claim that $\{\Sigma_t\}$ will be a foliation of $\mathbf{H}^3$. By Lemma 4.2, $\Sigma_t \cap \Sigma_s = \emptyset$ for any $s \neq t$. Hence, the only way to fail to be a foliation for $\{\Sigma_t\}$ is to have a gap between two leaves.

Now, assume that there is a gap between the leaves $\{\Sigma_t \mid t > s\}$ and $\{\Sigma_t \mid t < s\}$. This implies if we have sequences $\{\Gamma_{t_i^+}\}$ and $\{\Gamma_{t_i^-}\}$, where $t_i^+ \rightarrow s$ from positive side, and $t_i^- \rightarrow s$ from negative side, then one of the sequences $\{\Sigma_{t_i^+}\}$ and $\{\Sigma_{t_i^-}\}$ has no subsequences converging to Σ_s , because of the existence of the gap between the leaves (Recall that we assume there is a unique absolutely area minimizing surface Σ_t with $\partial_\infty \Sigma_t = \Gamma_t$). However, this contradicts to Lemma 2.5. So, $\{\Sigma_t\}$ will be a foliation of $\mathbf{H}^3$.

Now, by assumption, there are two distinct minimal surface M_1 and M_2 which are asymptotic to Γ_0 . This implies at least one of these surfaces is not a leaf of the foliation, and say M_2 , must intersect the leaves in the foliation nontrivially. Since $\{\Sigma_t\}$ foliates whole $\mathbf{H}^3$, this means M_2 must intersect tangentially (lying in one side) one of the leaves in the foliation. However, this contradicts to the maximum principle. So, one of the simple closed curves in $\{\Gamma_t\}$ must bound more than one absolutely area minimizing surface in $\mathbf{H}^3$. The proof follows. $\square$

By using similar ideas, one can prove an analogous theorem for least area planes in $\mathbf{H}^3$.

Theorem 5.3. *There exists a simple closed curve Γ in $S_\infty^2(\mathbf{H}^3)$ such that Γ bounds more than one least area plane $\{P_i\}$ in $\mathbf{H}^3$, i.e. $\partial_\infty P_i = \Gamma$.*

Proof: The proof is completely analogous to the proof of previous theorem. Again, we start with the same foliation of $S_\infty^2(\mathbf{H}^3)$ with simple closed curves $\{\Gamma_t\}$ containing Γ_0 which bounds more than one complete minimal surface in $\mathbf{H}^3$.

Assume that P_t is the unique least area plane in $\mathbf{H}^3$ with $\partial_\infty P_t = \Gamma_t$. Like in the proof of the theorem above, by using Lemma 3.1, and Lemma 4.2, it can be showed that the family of least area planes $\{P_t\}$ foliates $\mathbf{H}^3$. However, like in above theorem, Γ_0 bounds more than one area minimizing surface, and at least one of them is not a leaf of the foliation, say M_2 . Hence, M_2 must intersect tangentially (lying in one side) one of the leaves in the foliation. Again, this contradicts to the maximum principle for minimal surfaces. The proof follows. $\square$

Remark 5.2. The same proof may not work for area minimizing surfaces in a specified topological class. The problem is that Lemma 2.5 may not be true for this case as the limiting surface might not be in the same topological class.

Remark 5.3. As the introduction suggests, there is no known example of a simple closed curve of $S_\infty^2(\mathbf{H}^3)$ with nonunique solution to the asymptotic Plateau problem. Unfortunately, the results above show the existence of such an example, but they do not give one. The main problem to find such an example is the noncompactness of the objects. In compact case, the quantitative data enables you to find such examples like baseball curve on a sphere, but in the asymptotic case the techniques do not work because of the lack of the quantitative data. To overcome this problem, it might be possible to employ the renormalized area defined by Alexakis-Mazzeo in [AM] in order to construct an explicit simple closed curve in $S_\infty^2(\mathbf{H}^3)$ bounding more than one absolutely area minimizing surface.

6. CONCLUDING REMARKS

In this paper, we showed that the space of closed, codimension-1 submanifolds of $S_\infty^{n-1}(\mathbf{H}^n)$ has a dense (and generic in a sense) subspace of closed, codimension-1 submanifolds of $S_\infty^{n-1}(\mathbf{H}^n)$ bounding a unique absolutely area minimizing hypersurface in $\mathbf{H}^n$. As we discussed in the introduction, Anderson showed this result for closed submanifolds bounding convex domains in $S_\infty^{n-1}(\mathbf{H}^n)$ in [A1]. Then, Hardt and Lin generalized this result to closed submanifolds bounding star shaped domains in $S_\infty^{n-1}(\mathbf{H}^n)$ in [HL]. These were the only cases known so far. Hence, our result shows how abundant they are. In dimension 3, they are generic, and in the higher dimensions, they are dense.

The technique which we employ here is very general, and it applies to many different settings of Plateau problem. In particular, it can naturally be generalized to the Gromov-Hadamard spaces which is studied by Lang in [L], and it can be generalized to the mean convex domains with spherical boundary which is studied by Lin in [Li]. In other words, codimension-1 closed submanifolds in the boundary of these spaces generically bounds unique absolutely area minimizing hypersurfaces. Generalizing this technique in the context of constant mean curvature hypersurfaces in hyperbolic space also gives similar results. On the other hand, they can also be applied in Gromov hyperbolic 3-spaces with cocompact metric where the author solved the asymptotic Plateau problem [Co3].

On the other hand, it was not known whether all closed codimension-1 submanifolds in $S_\infty^{n-1}(\mathbf{H}^n)$ have a unique solution to the asymptotic Plateau problem or not. The only known results about nonuniqueness are also come from Anderson in [A2]. He constructs examples of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ bounding more than one complete *minimal* surface in $\mathbf{H}^3$. These examples are also area minimizing in their topological class. However, none of them are absolutely area minimizing, i.e. a solution to the asymptotic Plateau problem. In Section 5, we prove the existence of simple closed curves in $S_\infty^2(\mathbf{H}^3)$ with nonunique solution to asymptotic Plateau problem, and hence, give an answer for dimension 3. However, there is no result in higher dimensions yet. In other words, it is not known whether there exist closed codimension-1 submanifolds in $S_\infty^{n-1}(\mathbf{H}^n)$ bounding more than one absolutely area minimizing hypersurfaces or not for $n > 3$.

It might be possible to extend the techniques in this paper to address the nonuniqueness question in higher dimensions. It is possible to use the technique in Remark 5.1 (Figure 3), to get a codimension-1 sphere Γ_0 in $S_\infty^{n-1}(\mathbf{H}^n)$ bounding an absolutely area minimizing hypersurface Σ_0 in $\mathbf{H}^n$ which is not a hyperplane. Like in the proof of Theorem 5.2, by foliating $S_\infty^{n-1}(\mathbf{H}^n)$ by closed, codimension-1 submanifolds $\{\Gamma_t\}$, and by assuming uniqueness of absolutely area minimizing hypersurfaces, one can get a foliation of $\mathbf{H}^n$ by the absolutely area minimizing hypersurfaces $\{\Sigma_t\}$ with $\partial_\infty \Sigma_t = \Gamma_t$. Let γ be a $n - 2$ -sphere in Σ_0 which is "close" to Γ_0 . Then by using [Wh2], one can get a compact area minimizing hyperplane M whose boundary is γ . Hence by construction, M cannot be in a leaf in the foliation $\{\Sigma_t\}$. Like in the proof of Theorem 5.2, it might be possible to get a contradiction by studying the intersection of M with the foliation $\{\Sigma_t\}$.

Unfortunately, the maximum principle [Si] does not work here since while Σ_t is absolutely area minimizing, M is not. Also, the maximum principles due to Solomon-White [SW] and Ilmanen [I] which are for stationary varifolds are not enough to get a contradiction since the minimizing hyperplane M given by [Wh2] might have codimension-1 singularities, while

[SW] and [I] works up to codimension-2 singularities. So, to get a contradiction here, one needs a stronger maximum principle, or more regular area minimizing hyperplane M .

REFERENCES

- [A1] M. Anderson, *Complete minimal varieties in hyperbolic space*, Invent. Math. **69** (1982) 477–494.
- [A2] M. Anderson, *Complete minimal hypersurfaces in hyperbolic n -manifolds*, Comment. Math. Helv. **58** (1983) 264–290.
- [AM] S. Alexakis, R. Mazzeo, *Renormalized area and properly embedded minimal surfaces in hyperbolic 3-manifolds*, arXiv:0802.2250
- [Co1] B. Coskunuzer, *Minimal Planes in Hyperbolic Space*, Comm. Anal. Geom. **12** (2004) 821–836.
- [Co2] B. Coskunuzer, *Generic Uniqueness of Least Area Planes in Hyperbolic Space*, Geom. & Topology **10** (2006) 401–412.
- [Co3] B. Coskunuzer, *Uniform 1-cochains and Genuine Laminations*, Topology **45** (2006) 751–784.
- [Co4] B. Coskunuzer, *Mean Convex Hulls and Least Area Disks spanning Extreme Curves*, Math Z. **252** (2006) 811–824.
- [Co5] B. Coskunuzer, *Asymptotic Plateau problem*, eprint; arXiv:0907.0552
- [Fe] H. Federer, *Geometric measure theory*, Springer-Verlag, New York 1969.
- [Ga] D. Gabai, *On the geometric and topological rigidity of hyperbolic 3-manifolds*, J. Amer. Math. Soc. **10** (1997) 37–74.
- [HL] R. Hardt and F.H. Lin, *Regularity at infinity for absolutely area minimizing hypersurfaces in hyperbolic space*, Invent. Math. **88** (1987) 217–224.
- [I] T. Ilmanen, *A strong maximum principle for singular minimal hypersurfaces*, Calc. Var. Partial Differential Equations **4** (1996) 443–467.
- [L] U. Lang, *Asymptotic Plateau problem in Gromov hyperbolic manifolds*, Calc. Var. Partial Differential Equations **16** (2003) 31–46.
- [Li] F.H. Lin, *Plateau’s problem for H -convex curves*, Manuscripta Math. **58** (1987) 497–511.
- [MY] W. Meeks and S.T. Yau, *The classical Plateau problem and the topology of three manifolds*, Topology **21**
- [Si] L. Simon, *A strict maximum principle for area minimizing hypersurfaces*, J. Differential Geom. **26** (1987) 327–335.
- [SW] B. Solomon and B. White, *A strong maximum principle for varifolds that are stationary with respect to even parametric elliptic functionals*, Indiana Univ. Math. J. **38** (1989) 683–691.
- [Wh1] B. White, *On the topological type of minimal submanifolds*, Topology **31** (1992) 445–448.
- [Wh2] B. White, *Mappings that minimize area in their homotopy classes*, J. Differential Geom. **20** (1984) 433–446.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06520
 Current address: Koc University, Department of Mathematics, Istanbul, Turkey
 E-mail address: bcoskunuzer@ku.edu.tr